

# Equations with Discontinuous Right Hand Side

L. Fridman

UNAM

- 1 Preliminaries
  - Absolute Continuity
  - Upper semi-continuity
- 2 Equations with Discontinuous RHS
  - Historical Remarks
- 3 Regularization Procedure for ODE with Discontinuous RHS
  - Filippov Solutions
  - Utkin Solutions
  - Aizerman-Pyatniskii Solutions
- 4 Disturbed Systems and Extended Differential Inclusions
  - Disturbances and Differential Inclusions
- 5 Existence of Solutions
  - Existence Conditions

- 1 Preliminaries
  - Absolute Continuity
  - Upper semi-continuity
- 2 Equations with Discontinuous RHS
  - Historical Remarks
- 3 Regularization Procedure for ODE with Discontinuous RHS
  - Filippov Solutions
  - Utkin Solutions
  - Aizerman-Pyatniskii Solutions
- 4 Disturbed Systems and Extended Differential Inclusions
  - Disturbances and Differential Inclusions
- 5 Existence of Solutions
  - Existence Conditions

## Definition

Let  $\mathcal{I}$  be an interval in the real line  $\mathbb{R}$ . A function  $f : \mathcal{I} \rightarrow \mathbb{R}$  is absolutely continuous on  $\mathcal{I}$  if for every positive number  $\epsilon$ , there is a positive number  $\delta$  such that whenever a finite sequence of pairwise disjoint sub-intervals  $(x_k; y_k)$  of  $\mathcal{I}$  satisfies

$$\sum_k (y_k - x_k) < \delta$$

then

$$\sum_k |f(y_k) - f(x_k)| < \epsilon$$

The collection of all absolutely continuous functions on  $\mathcal{I}$  is denoted  $AC(\mathcal{I})$ .

# Absolute continuity of functions

## Equivalent Definitions

- 1  $f$  is absolutely continuous
- 2  $f$  has a Lebesgue integrable derivative  $f'$  almost everywhere and

$$f(x) = f(a) + \int_a^x f'(t)dt; \quad \forall x \in [a; b]$$

- 3 there exists a Lebesgue integrable function  $g$  on  $[a; b]$  such that

$$f(x) = f(a) + \int_a^x g(t)dt; \quad \forall x \in [a; b]$$

If these equivalent conditions are satisfied then necessarily  $g = f'$  almost everywhere. Equivalence between (1) and (3) is known as the fundamental theorem of Lebesgue integral calculus, due to Lebesgue.


## Properties

- 1 If  $f, g \in AC(\mathcal{I})$ , then  $f \pm g$  is absolutely continuous.
- 2 If  $\mathcal{I}$  is a bounded closed interval and  $f, g \in AC(\mathcal{I})$ , then  $fg$  is also absolutely continuous.
- 3 If  $\mathcal{I}$  is a bounded closed interval,  $f \in AC(\mathcal{I})$  and  $f \neq 0$  then  $\frac{1}{f}$  is absolutely continuous.
- 4 Every absolutely continuous function is uniformly continuous and, therefore, continuous. Every Lipschitz-continuous function is absolutely continuous.
- 5 If  $f : \mathcal{I} \rightarrow \mathbb{R}$  is absolutely continuous, then it is of bounded variation on  $[a; b]$ .

## Example

$$f(x) = \begin{cases} 0 & \text{if } x = 0 \\ x \sin\left(\frac{1}{x}\right) & \text{if } x \neq 0 \end{cases}$$


on a finite interval containing the origin.



## Example

$$f(x) = |x|^{1/2}$$

in zero it is not differentiable and the lateral derivatives do not exist.





- 1 Preliminaries
  - Absolute Continuity
  - Upper semi-continuity
- 2 Equations with Discontinuous RHS
  - Historical Remarks
- 3 Regularization Procedure for ODE with Discontinuous RHS
  - Filippov Solutions
  - Utkin Solutions
  - Aizerman-Pyatniskii Solutions
- 4 Disturbed Systems and Extended Differential Inclusions
  - Disturbances and Differential Inclusions
- 5 Existence of Solutions
  - Existence Conditions

# Upper semi-continuity of set-valued functions

Introduce the following distances

$$\rho(x, \mathbf{M}) = \inf_{y \in \mathbf{M}} \|x - y\|, \quad x \in \mathbb{R}^n, \mathbf{M} \subset \mathbb{R}^n,$$

$$\rho(\mathbf{M}_1, \mathbf{M}_2) = \sup_{x \in \mathbf{M}_1} \rho(x, \mathbf{M}_2), \quad \mathbf{M}_1 \subset \mathbb{R}^n, \mathbf{M}_2 \subset \mathbb{R}^n,$$

In general, the distance  $\rho$  is not symmetric,  $\rho(\mathbf{M}_1, \mathbf{M}_2) \neq \rho(\mathbf{M}_2, \mathbf{M}_1)$ .


# Upper semi-continuity of set-valued functions

Introduce the following distances

$$\rho(x, \mathbf{M}) = \inf_{y \in \mathbf{M}} \|x - y\|, \quad x \in \mathbb{R}^n, \mathbf{M} \subset \mathbb{R}^n,$$

$$\rho(\mathbf{M}_1, \mathbf{M}_2) = \sup_{x \in \mathbf{M}_1} \rho(x, \mathbf{M}_2), \quad \mathbf{M}_1 \subset \mathbb{R}^n, \mathbf{M}_2 \subset \mathbb{R}^n,$$

In general, the distance  $\rho$  is not symmetric,  $\rho(\mathbf{M}_1, \mathbf{M}_2) \neq \rho(\mathbf{M}_2, \mathbf{M}_1)$ .



## Definition


A set-valued function  $F : \mathbb{R}^{n+1} \rightarrow 2^{\mathbb{R}^{n+1}}$  is said to be **upper semi-continuous** at a point  $(t^*, x^*) \in \mathbb{R}^{n+1}$  if  $(t, x) \rightarrow (t^*, x^*)$  implies

$$\rho(F(t, x), F(t^*, x^*)) \rightarrow 0.$$

## Example (Upper semi-continuous set-valued function)

$$\overline{\text{sign}}[\rho] = \begin{cases} 1 & \text{if } \rho > 0 \\ -1 & \text{if } \rho < 0 \\ [-1, 1] & \text{if } \rho = 0 \end{cases}$$

is an upper semi-continuous set-valued function.



- 1 Preliminaries
  - Absolute Continuity
  - Upper semi-continuity
- 2 Equations with Discontinuous RHS
  - Historical Remarks
- 3 Regularization Procedure for ODE with Discontinuous RHS
  - Filippov Solutions
  - Utkin Solutions
  - Aizerman-Pyatniskii Solutions
- 4 Disturbed Systems and Extended Differential Inclusions
  - Disturbances and Differential Inclusions
- 5 Existence of Solutions
  - Existence Conditions

## Differential Equations with Discontinuous RHS

$$\dot{x}(t) = f(t, x(t)),$$
$$t \in \mathbb{R}, x \in \mathbb{R}^n, f : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$$

- RHS Discontinuous with respect to the **time** variable (Caratheodory 1927)




Constantin Caratheodory  
(1873-1950)

## Differential Inclusions (Contingent Differential Equations)

$$\dot{x}(t) \in F(t, x(t)),$$

$$t \in \mathbb{R}, x \in \mathbb{R}^n, F : \mathbb{R} \times \mathbb{R}^n \rightarrow 2^{\mathbb{R}^n}$$

(Zaremba 1936, Marchaud 1938,  
Filippov 1959, Wazawski 1961,  
Cellina 1970, Antosiewicz 1975,  
Tolstonogov 1981, Aubin 1984)



Stanislaw Zaremba  
(1863-1942)



## Differential Equations with Discontinuous RHS

$$\dot{x}(t) = f(t, x(t)), \quad t \in \mathbb{R}, \quad x \in \mathbb{R}^n, \quad f : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$$

- **RHS Discontinuous with respect to the state variable** (Filippov 1960, Utkin 1967, Aizerman & Pyatnitskii 1974)



Professors A. Filippov, E. Pyatnitskii, M. Aizerman and V. Utkin

- 1 Preliminaries
  - Absolute Continuity
  - Upper semi-continuity
- 2 Equations with Discontinuous RHS
  - Historical Remarks
- 3 Regularization Procedure for ODE with Discontinuous RHS
  - **Filippov Solutions**
  - Utkin Solutions
  - Aizerman-Pyatniskii Solutions
- 4 Disturbed Systems and Extended Differential Inclusions
  - Disturbances and Differential Inclusions
- 5 Existence of Solutions
  - Existence Conditions

# ODE with Discontinuous RHS

$$\dot{x}(t) = f(t, x(t)), \quad t \in \mathbb{R}, \quad x \in \mathbb{R}^n, \quad f : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n \quad (\text{DiscRHS})$$

# ODE with Discontinuous RHS


$$\dot{x}(t) = f(t, x(t)), \quad t \in \mathbb{R}, \quad x \in \mathbb{R}^n, \quad f : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n \quad (\text{DiscRHS})$$

$f$  is **piecewise continuous**:

- 1  $\mathbb{R}^{n+1} = \bigcup_{j=1}^N \bar{G}_j$ , where  $G_j$ -open  
connected sets  $G_i \cap G_j \neq \emptyset, i \neq j$ ;
- 2  $\mathcal{S} \bigcup_{j=1}^N \partial G_j$  is of measure zero;
- 3  $f(t, x)$  is continuous in each  $G_j$  and  
 $\forall (t, x) \in \partial G_j : \exists f^j(t, x) \in \mathbb{R}^n$

$$f^j = \lim_{(t^k, x^k) \rightarrow (t, x)} f(t^k, x^k),$$

$$(t^k, x^k) \in G_j, \quad (t, x) \in \partial G_j$$




$$\dot{x}(t) \in F(t, x(t)), \quad t \in \mathbb{R} \quad (\text{DiffInc})$$
$$F(t, x) = \begin{cases} \{f(t, x)\} & \text{if } (t, x) \in \mathbb{R}^{n+1} \setminus \mathcal{S}, \\ \text{co} \left( \bigcup_{j \in \mathcal{N}(t, x)} \{f^j(t, x)\} \right) & \text{if } (t, x) \in \mathcal{S}, \end{cases}$$
$$\mathcal{N}(t, x) = \{j \in \{1, 2, \dots, N\} : (t, x) \in \partial G_j\}.$$

$$\dot{x}(t) \in F(t, x(t)), \quad t \in \mathbb{R} \quad (\text{DiffInc})$$
$$F(t, x) = \begin{cases} \{f(t, x)\} & \text{if } (t, x) \in \mathbb{R}^{n+1} \setminus \mathcal{S}, \\ \text{co} \left( \bigcup_{j \in \mathcal{N}(t, x)} \{f^j(t, x)\} \right) & \text{if } (t, x) \in \mathcal{S}, \end{cases}$$
$$\mathcal{N}(t, x) = \{j \in \{1, 2, \dots, N\} : (t, x) \in \partial G_j\}.$$


## Definition (Filippov 1960)

An absolutely continuous function  $x : \mathcal{I} \rightarrow \mathbb{R}^n$  defined on some interval or segment  $\mathcal{I}$  is called a solution of (DiscRHS) if it satisfies the differential inclusion (DiffInc) almost everywhere on  $\mathcal{I}$ .


# Illustration of Filippov regularization




(a) Switching case.




(b) Sliding mode.




# Illustration of Filippov regularization



(a) Switching case.



(b) Sliding mode.



## Example


$$\dot{x}(t) = -\text{sign}[x(t)] + d(t), t > 0,$$

where  $x(\cdot) \in \mathbb{R}$ ,  $\|d\|_C \leq d_0 < 1$ .


$$\text{sign}[\rho] = \begin{cases} 1 & \text{if } \rho > 0 \\ -1 & \text{if } \rho < 0 \\ 0 & \text{if } \rho = 0 \end{cases}$$




# Illustration of Filippov regularization



(a) Switching case.



(b) Sliding mode.



## Example

$$\dot{x}(t) \in -\overline{\text{sign}}[x(t)] + d(t), t > 0,$$

where  $x(\cdot) \in \mathbb{R}$ ,  $\|d\|_{\mathbb{C}} \leq d_0 < 1$ .

$$\overline{\text{sign}}[\rho] = \begin{cases} 1 & \text{if } \rho > 0 \\ -1 & \text{if } \rho < 0 \\ [-1, 1] & \text{if } \rho = 0 \end{cases}$$

- 1 Preliminaries
  - Absolute Continuity
  - Upper semi-continuity
- 2 Equations with Discontinuous RHS
  - Historical Remarks
- 3 Regularization Procedure for ODE with Discontinuous RHS
  - Filippov Solutions
  - **Utkin Solutions**
  - Aizerman-Pyatniskii Solutions
- 4 Disturbed Systems and Extended Differential Inclusions
  - Disturbances and Differential Inclusions
- 5 Existence of Solutions
  - Existence Conditions

# Discontinuous Control Systems

Let us consider the system

$$\dot{x}(t) = f(t, x(t), u(t, x(t))), t \in \mathbb{R}, \quad (\text{DisContSys})$$

where  $f : \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^n$ ,  $f \in \mathbb{C}$  and

$$u : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^m, \quad u(t, x) = (u_1(t, x), u_2(t, x), \dots, u_m(t, x))^T$$

is a **piecewise continuous** feedback control

# Discontinuous Control Systems

Let us consider the system

$$\dot{x}(t) = f(t, x(t), u(t, x(t))), t \in \mathbb{R}, \quad (\text{DisContSys})$$

where  $f : \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^n$ ,  $f \in \mathbb{C}$  and

$$u : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^m, \quad u(t, x) = (u_1(t, x), u_2(t, x), \dots, u_m(t, x))^T$$

is a **piecewise continuous** feedback control

## Assumption

Each component  $u_i(t, x)$  is discontinuous only on a surface

$$\mathcal{S}_i = \{(t, x) \in \mathbb{R}^n : s_i(t, x) = 0\},$$

where functions  $s_i : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  are smooth, i.e.  $s_i \in \mathbb{C}^1(\mathbb{R}^{n+1})$ .

# Utkin Regularization

$$\dot{x}(t) = f(t, x(t), U(t, x(t))), t \in \mathbb{R},$$

where  $U(t, x) = (U_1(t, x), U_2(t, x), \dots, U_m(t, x))^T$  and

$$U_i(t, x) = \begin{cases} u_i(t, x), & s_i(t, x) \neq 0 \\ \text{co} \left\{ \lim_{\substack{(t_j, x_j) \rightarrow (t, x) \\ s_i(t_j, x_j) > 0}} u_i(t_j, x_j), \lim_{\substack{(t_j, x_j) \rightarrow (t, x) \\ s_i(t_j, x_j) < 0}} u_i(t_j, x_j) \right\}, & s_i(t, x) = 0 \end{cases},$$

(ValFunc)

The set  $f(t, x, U(t, x))$  is **non-convex** in general case.

## Example (Utkin Regularization)

$$u(x) = -\text{sign}[x] \quad \text{and} \quad U(x) = \overline{\text{sign}}[x]$$

$$\text{sign}[\rho] = \begin{cases} 1 & \text{if } \rho > 0 \\ -1 & \text{if } \rho < 0 \\ 0 & \text{if } \rho = 0 \end{cases}, \quad \overline{\text{sign}}[\rho] = \begin{cases} 1 & \text{if } \rho > 0 \\ -1 & \text{if } \rho < 0 \\ [-1, 1] & \text{if } \rho = 0 \end{cases}$$

# Equivalent Control (Utkin Solution)


## Definition

An absolutely continuous function  $x : \mathcal{I} \rightarrow \mathbb{R}^n$  defined on some interval or segment  $\mathcal{I}$  is called a solution of (DisContSys) if there exists a measurable function  $u_{eq} : \mathcal{I} \rightarrow \mathbb{R}^m$  such that  $u_{eq}(t) \in U(t, x(t))$  and  $\dot{x}(t) = f(t, x(t), u_{eq}(t))$  almost everywhere on  $\mathcal{I}$ .


# Equivalent Control (Utkin Solution)

## Definition

An absolutely continuous function  $x : \mathcal{I} \rightarrow \mathbb{R}^n$  defined on some interval or segment  $\mathcal{I}$  is called a solution of (DisContSys) if there exists a measurable function  $u_{eq} : \mathcal{I} \rightarrow \mathbb{R}^m$  such that  $u_{eq}(t) \in U(t, x(t))$  and  $\dot{x}(t) = f(t, x(t), u_{eq}(t))$  almost everywhere on  $\mathcal{I}$ .



(a) Filippov definition.



(b) Utkin definition.

Equivalent control (Utkin 1967):  $s(x) = 0$  and  $\frac{\partial s(x)}{\partial x} f(t, x, u_{eq}) = 0$



## Example (Equivalent Control)

$$\dot{x}_1 = u$$

$$\dot{x}_2 = (2u^2 - 1)x_2$$

$$u(t) = -\text{sign}[x_1(t)]$$

## Example (Equivalent Control)

$$\dot{x}_1 = u$$

$$\dot{x}_2 = (2u^2 - 1)x_2$$

$$u(t) = -\text{sign}[x_1(t)]$$

Filippov definition

$$\begin{bmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{bmatrix} \in \begin{bmatrix} -\overline{\text{sign}}[x_1(t)] \\ x_2(t) \end{bmatrix}$$

## Example (Equivalent Control)

$$\dot{x}_1 = u$$

$$\dot{x}_2 = (2u^2 - 1)x_2$$

$$u(t) = -\text{sign}[x_1(t)]$$

Filippov definition

$$\begin{bmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{bmatrix} \in \begin{bmatrix} -\overline{\text{sign}}[x_1(t)] \\ x_2(t) \end{bmatrix}$$

**Unstable**

## Example (Equivalent Control)

$$\dot{x}_1 = u$$

$$\dot{x}_2 = (2u^2 - 1)x_2$$

$$u(t) = -\text{sign}[x_1(t)]$$

Filippov definition

$$\begin{bmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{bmatrix} \in \begin{bmatrix} -\overline{\text{sign}}[x_1(t)] \\ x_2(t) \end{bmatrix}$$

Utkin definition

$$\begin{bmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{bmatrix} \in \begin{bmatrix} -\overline{\text{sign}}[x_1(t)] \\ -x_2(t) \end{bmatrix}$$

**Unstable**

## Example (Equivalent Control)

$$\dot{x}_1 = u$$

$$\dot{x}_2 = (2u^2 - 1)x_2$$

$$u(t) = -\text{sign}[x_1(t)]$$

Filippov definition

$$\begin{bmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{bmatrix} \in \begin{bmatrix} -\overline{\text{sign}}[x_1(t)] \\ x_2(t) \end{bmatrix}$$

**Unstable**

Utkin definition

$$\begin{bmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{bmatrix} \in \begin{bmatrix} -\overline{\text{sign}}[x_1(t)] \\ -x_2(t) \end{bmatrix}$$

**Stable**

## Example (Utkin 1970s)

$$\dot{x}_1 = 0.3x_2(t) + x_1(t)u(t),$$

$$\dot{x}_2 = -0.7x_1(t) + 4x_1^3(t)u(t),$$


$$u(t) = -\text{sign}[x_1(t)s(t)],$$

$$s(t) = x_1(t) + x_2(t),$$

## Example (Utkin 1970s)

$$\begin{aligned}\dot{x}_1 &= 0.3x_2(t) + x_1(t)u(t), \\ \dot{x}_2 &= -0.7x_1(t) + 4x_1^3(t)u(t),\end{aligned}$$

$$\begin{aligned}u(t) &= -\text{sign}[x_1(t)s(t)], \\ s(t) &= x_1(t) + x_2(t),\end{aligned}$$



- 1 Preliminaries
  - Absolute Continuity
  - Upper semi-continuity
- 2 Equations with Discontinuous RHS
  - Historical Remarks
- 3 Regularization Procedure for ODE with Discontinuous RHS
  - Filippov Solutions
  - Utkin Solutions
  - Aizerman-Pyatniskii Solutions
- 4 Disturbed Systems and Extended Differential Inclusions
  - Disturbances and Differential Inclusions
- 5 Existence of Solutions
  - Existence Conditions



# Aizerman-Pyatniskii Regularization (Filippov 1988)

$$\dot{x} \in \text{co}(f(t, x, U(t, x))), \quad t \in \mathbb{R}$$

## Definition


An absolutely continuous function  $x : \mathcal{I} \rightarrow \mathbb{R}^n$  defined on some interval or segment  $\mathcal{I}$  is called a solution of (DiscRHS) if it satisfies the differential inclusion (DiffInc) almost everywhere on  $\mathcal{I}$ .

# Aizerman-Pyatniskii Regularization (Filippov 1988)


$$\dot{x} \in \text{co}(f(t, x, U(t, x))), \quad t \in \mathbb{R}$$

## Definition


An absolutely continuous function  $x : \mathcal{I} \rightarrow \mathbb{R}^n$  defined on some interval or segment  $\mathcal{I}$  is called a solution of (DiscRHS) if it satisfies the differential inclusion (DiffInc) almost everywhere on  $\mathcal{I}$ .



(a) Filippov



(b) Utkin



(c) Aizerman-Pyatniskii

## Presence of fast actuators (Fridman 2001,2002)

### Actuators

1st order actuator

$$\mu \dot{z}_1 = -2z_1 - u(s),$$

2nd order actuator

$$\begin{aligned}\mu \dot{z}_1 &= z_2, \\ \mu \dot{z}_2 &= -2z_1 - 3z_2 - u(s),\end{aligned}$$

### Plant

$$\dot{s} = z, \quad \dot{x} = u^4 - u^2 + \beta x, \quad u(s) = \text{sign}[s],$$

## Presence of fast actuators (Fridman 2001,2002)

### Actuators

1st order actuator

$$\mu \dot{z}_1 = -2z_1 - u(s),$$

2nd order actuator

$$\begin{aligned}\mu \dot{z}_1 &= z_2, \\ \mu \dot{z}_2 &= -2z_1 - 3z_2 - u(s),\end{aligned}$$

### Plant

$$\dot{s} = z, \quad \dot{x} = u^4 - u^2 + \beta x, \quad u(s) = \text{sign}[s],$$

## Presence of fast actuators (Fridman 2001,2002)

### Actuators

1st order actuator

$$\mu \dot{z}_1 = -2z_1 - u(s),$$

2nd order actuator

$$\begin{aligned}\mu \dot{z}_1 &= z_2, \\ \mu \dot{z}_2 &= -2z_1 - 3z_2 - u(s),\end{aligned}$$

### Plant

$$\dot{s} = z, \quad \dot{x} = u^4 - u^2 + \beta x, \quad u(s) = \text{sign}[s],$$

## Presence of fast actuators (Fridman 2001,2002)

### Actuators

1st order actuator

$$\mu \dot{z}_1 = -2z_1 - u(s),$$

2nd order actuator

$$\begin{aligned}\mu \dot{z}_1 &= z_2, \\ \mu \dot{z}_2 &= -2z_1 - 3z_2 - u(s),\end{aligned}$$

### Plant

$$\dot{s} = z, \quad \dot{x} = u^4 - u^2 + \beta x, \quad u(s) = \text{sign}[s],$$

## Presence of fast actuators (Fridman 2001,2002)

### Actuators

1st order actuator

$$\mu \dot{z}_1 = -2z_1 - u(s),$$

2nd order actuator

$$\begin{aligned}\mu \dot{z}_1 &= z_2, \\ \mu \dot{z}_2 &= -2z_1 - 3z_2 - u(s),\end{aligned}$$

### Plant

$$\dot{s} = z, \quad \dot{x} = u^4 - u^2 + \beta x, \quad u(s) = \text{sign}[s],$$

## Reduced Order System

$$\mu = 0 \Rightarrow z_1 = -u(s)/2, \quad \dot{s} = -u/2, \quad \dot{x} = (u^4 - u^2 + \beta)x,$$



## Reduced Order System

$$\mu = 0 \Rightarrow z_1 = -u(s)/2, \quad \dot{s} = -u/2, \quad \dot{x} = (u^4 - u^2 + \beta)x,$$

Sliding Dynamics (**Filippov=Utkin**)

1 st order actuator  $(z(t), s(t)) \rightarrow 0$  Sliding dynamics  $\dot{x} = \beta x \Rightarrow$  Unstable

2nd order actuator

$\exists \left( z_0 \left( \frac{t}{\mu} \right), s_0 \left( \frac{t}{\mu} \right) \right)$  – Periodic Solution

$\exists \bar{\beta}(\mu) : \forall \beta < \bar{\beta}(\mu) \exists \gamma :$

$$-\gamma = \int_0^T [(2z_1(\tau))^4 - (2z_2(\tau))^2] d\tau$$

$\Rightarrow \dot{x} = -(\gamma - \beta)x$

Could be stable

# Equivalence of Definitions

## Theorem (Utkin 1992, Zolezzi 2002)

Let a right-hand side of the system (DiscRHS) be affine with respect to control:

$$f(t, x, u(t, x)) = a(t, x) + b(t, x)u(t, x),$$

where  $a : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ ,  $b : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n \times m}$ ,  $a, b \in \mathbb{C}$  and  $u : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^m$  is a piecewise continuous function  $u(t, x) = (u_1(t, x), \dots, u_m(t, x))^T$ , such that  $u_i$  has a unique switching surface  $s_i(x) = 0$ ,  $s_i \in \mathbb{C}^1(\mathbb{R}^n)$ .

# Equivalence of Definitions

## Theorem (Utkin 1992, Zolezzi 2002)

Let a right-hand side of the system (DiscRHS) be affine with respect to control:

$$f(t, x, u(t, x)) = a(t, x) + b(t, x)u(t, x),$$

where  $a : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ ,  $b : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n \times m}$ ,  $a, b \in \mathbb{C}$  and  $u : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^m$  is a piecewise continuous function  $u(t, x) = (u_1(t, x), \dots, u_m(t, x))^T$ , such that  $u_i$  has a unique switching surface  $s_i(x) = 0$ ,  $s_i \in \mathbb{C}^1(\mathbb{R}^n)$ .

Definitions of Filippov, Utkin and Aizerman-Pyatnitskii are equivalent iff

$$\det \left( \nabla^T s(x) b(t, x) \right) \neq 0 \text{ if } (t, x) \in S,$$

where  $s(x) = (s_1(x), s_2(x), \dots, s_m(x))^T$ ,  $\nabla s(x) \in \mathbb{R}^{n \times m}$  is the matrix of partial derivatives  $\frac{\partial s_i}{\partial x_j}$  and  $S$  is a discontinuity set of  $u(t, x)$ .

## Example (Neimark 1961)

$$\begin{aligned}\dot{x} &= Ax(t) + cu_1(t) + bu_2(t), \\ t > 0, \quad x(\cdot) &= (x_1(\cdot), x_2(\cdot))^T \in \mathbb{R}^2, \\ A \in \mathbb{R}^{2 \times 2}, \quad b &= (0, 1)^T,\end{aligned}$$

$$\begin{aligned}u_1(t) &= -\text{sign}[x_1(t)], \\ u_2(t) &= -\text{sign}[x_1(t)], \\ c &= (1, 0)^T,\end{aligned}$$


## Example (Neimark 1961)

$$\begin{aligned}\dot{x} &= Ax(t) + cu_1(t) + bu_2(t), \\ t > 0, \quad x(\cdot) &= (x_1(\cdot), x_2(\cdot))^T \in \mathbb{R}^2, \\ A \in \mathbb{R}^{2 \times 2}, \quad b &= (0, 1)^T,\end{aligned}$$

$$\begin{aligned}u_1(t) &= -\text{sign}[x_1(t)], \\ u_2(t) &= -\text{sign}[x_1(t)], \\ c &= (1, 0)^T,\end{aligned}$$


### Filippov definition

$$\dot{x} \in \{Ax\} \dot{+} (b + c) \cdot \overline{\text{sign}[x_1]}$$



### Utkin definition

$$\dot{x} \in \{Ax\} \dot{+} b \cdot \overline{\text{sign}[x_1]} + c \cdot \overline{\text{sign}[x_1]}.$$



- 1 Preliminaries
  - Absolute Continuity
  - Upper semi-continuity
- 2 Equations with Discontinuous RHS
  - Historical Remarks
- 3 Regularization Procedure for ODE with Discontinuous RHS
  - Filippov Solutions
  - Utkin Solutions
  - Aizerman-Pyatniskii Solutions
- 4 Disturbed Systems and Extended Differential Inclusions
  - Disturbances and Differential Inclusions
- 5 Existence of Solutions
  - Existence Conditions

# Disturbances and Differential Inclusions

Models of sliding mode control systems usually have the form

$$\dot{x}(t) = f(t, x(t), u(t, x(t)), d(t)), \quad t \in \mathbb{R},$$

- $x(\cdot) \in \mathbb{R}^n$  is the vector of system states,
- $u(\cdot, \cdot) \in \mathbb{R}^m$  is the vector of control inputs,
- $d(\cdot) \in \mathbb{R}^k$  is the vector of disturbances,
- the function  $f : \mathbb{R}^{n+m+k+1} \rightarrow \mathbb{R}^n$  is assumed to be continuous,
- the control function  $u : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^m$  is piecewise continuous,
- the vector-valued function  $d : \mathbb{R} \rightarrow \mathbb{R}^k$  is assumed to be locally measurable and bounded as follows:

$$d_i^{\min} \leq d_i(t) \leq d_i^{\max}$$

where  $d(t) = (d_1(t), d_2(t), \dots, d_k(t))^T$ ,  $t \in \mathbb{R}$ .



## Example (Disturbed sliding mode system)

Consider the simplest disturbed sliding mode system

$$\dot{x}(t) = -d_1(t) \operatorname{sign}[x(t)] + d_2(t), \quad (\text{Ex1})$$

where  $x \in \mathbb{R}$ , unknown functions  $d_i : \mathbb{R} \rightarrow \mathbb{R}$  are bounded by

$$d_i^{\min} \leq d_i(t) \leq d_i^{\max}, \quad i = 1, 2.$$

## Example (Disturbed sliding mode system)

Consider the simplest disturbed sliding mode system

$$\dot{x}(t) = -d_1(t) \operatorname{sign}[x(t)] + d_2(t), \quad (\text{Ex1})$$

where  $x \in \mathbb{R}$ , unknown functions  $d_i : \mathbb{R} \rightarrow \mathbb{R}$  are bounded by

$$d_i^{\min} \leq d_i(t) \leq d_i^{\max}, \quad i = 1, 2.$$

Obviously, all solutions of the system (Ex1) belong to a solution set of the following extended differential inclusion

$$\dot{x}(t) \in - [d_1^{\min}, d_1^{\max}] \cdot \overline{\operatorname{sign}}[x(t)] + [d_2^{\min}, d_2^{\max}]. \quad (\text{Ex2})$$

Stability of the system (Ex2) implies the same property for (Ex1). In particular, for  $d_1^{\min} > \max\{|d_2^{\min}|, |d_2^{\max}|\}$  both these systems have **asymptotically stable origin**.

## Extended Differential Inclusion

All further considerations deal with the extended differential inclusion

$$\dot{x}(t) \in F(t, x(t)), \quad t \in \mathbb{R},$$

where

$$F(t, x) = \text{co}\{f(t, x, U(t, x), D)\},$$

the set-valued function  $U(t, x)$  is defined by (ValFunc) and

$$D = \begin{pmatrix} [d_1^{\min}, d_1^{\max}] \\ [d_2^{\min}, d_2^{\max}] \\ \vdots \\ [d_k^{\min}, d_k^{\max}] \end{pmatrix}$$

# Outline

- 1 Preliminaries
  - Absolute Continuity
  - Upper semi-continuity
- 2 Equations with Discontinuous RHS
  - Historical Remarks
- 3 Regularization Procedure for ODE with Discontinuous RHS
  - Filippov Solutions
  - Utkin Solutions
  - Aizerman-Pyatniskii Solutions
- 4 Disturbed Systems and Extended Differential Inclusions
  - Disturbances and Differential Inclusions
- 5 Existence of Solutions
  - Existence Conditions

## Theorem (Filippov 1960)

Let

- $F : \mathbf{G} \rightarrow 2^{\mathbb{R}^n}$  be **upper semi-continuous** at each point of the set

$$\mathbf{G} = \{(t, x) \in \mathbb{R}^{n+1} : |t - t_0| \leq a \text{ and } \|x - x_0\| < b,$$

where  $a, b \in \mathbb{R}_+$ ,  $t_0 \in \mathbb{R}$ ,  $x_0 \in \mathbb{R}^n$ ;

- $F(t, x)$  be **nonempty, compact and convex** for  $(t, x) \in \mathbf{G}$ ;
- there exists  $K > 0$  such that  $\rho(0, F(t, x)) < K$  for  $(t, x) \in \mathbf{G}$ ;

## Theorem (Filippov 1960)

Let

- $F : \mathbf{G} \rightarrow 2^{\mathbb{R}^n}$  be **upper semi-continuous** at each point of the set

$$\mathbf{G} = \{(t, x) \in \mathbb{R}^{n+1} : |t - t_0| \leq a \text{ and } \|x - x_0\| < b,$$

where  $a, b \in \mathbb{R}_+$ ,  $t_0 \in \mathbb{R}$ ,  $x_0 \in \mathbb{R}^n$ ;

- $F(t, x)$  be **nonempty, compact and convex** for  $(t, x) \in \mathbf{G}$ ;
- there exists  $K > 0$  such that  $\rho(0, F(t, x)) < K$  for  $(t, x) \in \mathbf{G}$ ;

**then**  $\exists x : \mathbb{R} \rightarrow \mathbb{R}^n$  - **absolutely continuous** and defined at least on  $[t_0 - \alpha, t_0 + \alpha]$ ,  $\alpha = \min\{a, b/K\}$ , such that  $x(t_0) = x_0$  and the inclusion

$$\dot{x}(t) \in F(t, x(t))$$

holds almost everywhere on  $[t_0 - \alpha, t_0 + \alpha]$ .

## Lemma (Filippov 1959)

Let

- a function  $f : \mathbb{R}^{n+m+1} \rightarrow \mathbb{R}^n$  be continuous;
- a set-valued function  $U : \mathbb{R}^{n+1} \rightarrow 2^{\mathbb{R}^m}$  be defined and upper-semicontinuous on an open set  $\mathcal{I} \times \Omega$ , where  $\Omega \subseteq \mathbb{R}^n$ ;
- $U(t, x)$  be nonempty, compact and convex for every  $(t, x) \in \mathcal{I} \times \Omega$ .
- a function  $x : \mathbb{R} \rightarrow \mathbb{R}^n$  be absolutely continuous on  $\mathcal{I}$ ,  $x(t) \in \Omega$  for  $t \in \mathcal{I}$ ,
- $\dot{x}(t) \in f(t, x(t), U(t, x(t)))$  almost everywhere on  $\mathcal{I}$ ;

# On existence of Utkin Solutions

## Lemma (Filippov 1959)

Let

- a function  $f : \mathbb{R}^{n+m+1} \rightarrow \mathbb{R}^n$  be continuous;
- a set-valued function  $U : \mathbb{R}^{n+1} \rightarrow 2^{\mathbb{R}^m}$  be defined and upper-semicontinuous on an open set  $\mathcal{I} \times \Omega$ , where  $\Omega \subseteq \mathbb{R}^n$ ;
- $U(t, x)$  be nonempty, compact and convex for every  $(t, x) \in \mathcal{I} \times \Omega$ .
- a function  $x : \mathbb{R} \rightarrow \mathbb{R}^n$  be absolutely continuous on  $\mathcal{I}$ ,  $x(t) \in \Omega$  for  $t \in \mathcal{I}$ ,
- $\dot{x}(t) \in f(t, x(t), U(t, x(t)))$  almost everywhere on  $\mathcal{I}$ ;

Then there exists a measurable function  $u_{eq} : \mathbb{R} \rightarrow \mathbb{R}^m$  such that

$$u_{eq}(t) \in U(t, x(t)) \quad \text{and} \quad \dot{x}(t) = f(t, x(t), u_{eq}(t))$$

almost everywhere on  $\mathcal{I}$ .



## Theorem (Gel'fand et al. 1978)

Let a set-valued function  $F : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$  be defined and upper-semicontinuous in  $\mathbb{R}^{n+1}$ .

Let  $F(t, x)$  be nonempty, compact and convex for any  $(t, x) \in \mathbb{R}^{n+1}$ .

If there exists a real valued function  $L : \mathbb{R}_+ \cup \{0\} \rightarrow \mathbb{R}_+ \cup \{0\}$  such that

$$\rho(0, F(t, x)) \leq L(\|x\|) \quad \text{and} \quad \int_0^{+\infty} \frac{1}{L(r)} dr = +\infty,$$

then for any  $(t_0, x_0) \in \mathbb{R}^{n+1}$  the system (DiffInc) has a solution  $x(t) : x(t_0) = x_0$  defined for all  $t \in \mathbb{R}$ .

## Summary

- Stability property of ODE with discontinuous RHS depends on definition of a solution.
- Stability of Aizerman-Pyatnitskii solutions always implies stability of Filippov and Utkin solutions.
- All introduced definitions may be equivalent in the case of affine control systems with discontinuous input.
- Analysis of the disturbed systems can be reduced to differential inclusions.